

Stefancik, Igor (2004): Tree species growth in mixed spruce, fir and beech stands; Folia Oecologica- vol. 31, no. 1.(2004): 36-39

Reference

Stefancik, Igor (2004): Tree species growth in mixed spruce, fir and beech stands; Folia Oecologica- vol. 31, no. 1.(2004): 36-39

Short reference

Stefancik (2004)

First author

Stefancik, Igor

Year

2004

Abstract

The paper presents a comparison of tree species growth on four research plots in Slovakia.

The plots are situated in mixed spruce, fir and beech stands where no interventions out in the last 30 years. The results showed that in the initial stage of research had fir the second highest proportion in tree species composition over three of the plots and/or on one of them even the highest one. After 30 years, the highest proportion of beech was found over the plots overall and/or the following total decrease (according to the number of trees) was registered: 76-89% for fir 46-93% for spruce and 36-73% for beech. After 30 years, beech reached the upper crown layer, while fir together with spruce constituted the medium crown layer. As for mean crown width, differences between individual tree species were low and statistically insignificant ($P > 0.05$). The highest mean annual diameter increment was registered for spruce and the lowest for fir.

[habitat: coniferous mixed woodlands](#)

[forest dynamic, gap dynamic, succession](#)

Journal

Folia Oecologica

Location

ER Archívum (2004/P-023/1, 2004/P-023/2)

Type

scientific paper

Attached document

[Tree species growth in mixed spruce, fir and beech stands](#) 227.3 KB

Katalógusba vette

Kovács Gabriella

Katalógusbavétel időpontja

Wed, 07/29/2015 - 12:00