

**Kanalas, P. et al. (2010): Seasonal and diurnal variability in sap flow intensity of mature sessile oak (Quercus petraea (Matt.) Liebl.) trees in relation to microclimatic conditions. Acta Biologica Hungarica 61: 95-108.**

Teljes hivatkozás

Kanalas, P. et al. (2010): Seasonal and diurnal variability in sap flow intensity of mature sessile oak (*Quercus petraea* (Matt.) Liebl.) trees in relation to microclimatic conditions. *Acta Biologica Hungarica* 61: 95-108.

Rövid hivatkozás

Kanalas (2010)

Első szerző

Kanalas P.

Év

2010

Összefoglalás

Seasonal and diurnal variability in sap flow intensity of mature sessile oak (*Quercus petraea* (Matt.) Liebl.) trees in relation to microclimatic conditions

P. Kanalas, A. Fenyvesi, J. Kis, E. Szöllősi, V. Oláh, I. Ander & I. Mészáros

In this study sap flow dynamics of mature sessile oak trees (*Quercus petraea*) in a marginal sessile oak-turkey oak forest was investigated in 2009. That year spring was dry without significant rain in April and May and the driest month was August. Due to the extreme weather conditions the volumetric soil water content (SWC) of upper 30 cm was low on experimental days in May (0.13-0.14 cm<sup>3</sup> cm<sup>-3</sup>) but it reached the lowest value in August (0.08 cm<sup>3</sup> cm<sup>-3</sup>). Sap flow was measured in a dominant and a co-dominant tree by heat dissipation method from 26 March till 30 October. In the present paper several three-day long periods of the continuous seasonal recordings were chosen to represent the effects of typical weather conditions and different stages of canopy development on sap flow dynamics. The daily maximum sap flow density values of dominant and co-dominant trees were similar (0.30-0.32 cm<sup>3</sup> cm<sup>-2</sup> min<sup>-1</sup>) in moist period (July). Rains and transient increase of SWC after proceeding drought resulted in change of diurnal course of sap flow in experimental days of July. In this period dominant trees also showed considerable sap flow (0.19 cm<sup>3</sup> cm<sup>-2</sup> min<sup>-1</sup>) during night

hours and short sap flow peaks in early morning (6:00 to 8:00 a.m.) indicating the refilling of desiccated tissues. After the progressive drought in August the daily maximum sap flow density decreased to 0.07 cm<sup>3</sup> cm<sup>-2</sup> min<sup>-1</sup> in dominant tree and to 0.12 cm<sup>3</sup> cm<sup>-2</sup> min<sup>-1</sup> in the co-dominant. Both trees exhibited gradual stomatal closure from morning hours.

[élőhely: fényben gazdag tölgyesek](#)

[erdőökológia](#)

[klíma: mikroklíma, állományklíma](#)

Megjegyzések

Drought stress, marginal forest, sap flow, sessile oak, stomatal control

Címszavazva - GE

Kiadó

Akadémiai Kiadó

Folyóirat

Acta Biologica Hungarica

Lelőhely

ER Archívum (2010/P-033)

Típus

tudományos folyóiratcikk

Katalógusba vette

Gulyás Györgyi

Katalógusbavétel időpontja

k, 07/05/2011 - 12:00